

CRISIS CENTER FOLLOW UP TO SAVE RESOURCES AND SAVE LIVES

Suicide is a national public health crisis, and is the tenth leading cause of death in the United States, with over 38,000 reported lives lost to suicide in the US in 2010. (Centers for Disease Control and Prevention (CDC), 2014). Research indicates that follow-up with hotline callers and people recently discharged from an emergency department (ED) or inpatient setting has positive results for both consumers and providers of mental health services. Follow-up has been shown to be cost effective and prevent suicides and crisis centers are uniquely positioned to be a crucial resource for people in need of follow-up care.

WHAT IS FOLLOW-UP?

Follow-up care can involve home visits, letters, phone calls, emails, or texts that are designed to check in with individuals who have recently experienced a suicide crisis to assess their well-being and level of risk and to support them as they continue their journey towards recovery. Follow-up is usually by telephone and typically occurs between 24 – 48 hours after the initial contact. Phone calls are brief and while they can be tailored to the individuals need, they are structured and focus on continued assessment of risk.

WHY IS FOLLOW-UP IMPORTANT?

Follow-up ensures continuity of care, provides support during a time of heightened risk, and facilitates linkages to outpatient care. It fills the significant gap and acts as a safety net for those at risk of suicide.

- Suicide risk is highest in the first week after discharge from an inpatient setting. This risk is 102 times higher in men and 246 times higher in women than their counterparts in the general population (Qin & Nordentoft, 2005).
- As many as 70% of suicide attempters never attend their first appointment or maintain treatment for more than a few sessions (multiple references).

BENEFITS OF FOLLOW-UP: SAVING LIVES

- Use of 24 hour crisis teams and 7 day follow-up programs show a significant reduction in suicide within 3 months of a patient's discharge from inpatient services (While et al., 2012).
- A study based in five countries indicated that follow up after emergency department discharge significantly reduced suicide. The follow-up program included 9 contacts by trained professionals at crisis centers over a maximum period of 18 months (Fleischmann, 2008).
- Patients who receive follow-up have a lower suicide rate in five years and a significantly lower suicide rate in the first two years after discharge (Motto & Bostrom, 2001).
- Following up with patients by telephone within one month after an emergency department discharge for a suicide attempt significantly reduces the likelihood that the person re-attempts suicide (Vaiva et al., 2006).
- In Australia, a study indicated that proactive telephone support for individuals with recurrent admissions reduced the number of hospital days per patient by 45% and saved \$AU895 per person (Andrews & Sunderland, 2009).
- Further, telephonic follow up before a service appointment results in improved motivation, a reduction in barriers to accessing services, and higher attendance rates (Zanjani, Miller, Turiano, Ross, & Oslin, 2008).
- In one year, a crisis center in St. Louis, Missouri reduced psychiatric hospitalization state-wide by 7% by referring some callers to more appropriate mobile outreach services and outpatient facilities based on the callers' needs (National Suicide Prevention Lifeline, 2011).

BENEFITS OF FOLLOW-UP: SAVING RESOURCES

- 45% of incurred costs for suicide attempt admissions are a result of readmissions to the ED (Beautrais & Gibbs, 2004).
- In a study of the return on investment (ROI) of post-discharge follow-up calls for suicidal ideation or deliberate self-harm, Truven Health Analytics estimated what the cost savings could be if an investment was made in crisis centers to place the follow-up calls. Truven estimated the amount of savings by reviewing data from crisis centers' cost of implementing follow-up calls and their potential to reduce hospital readmissions and additional emergency department visits within 30 days of discharge. The conclusion of this ROI analysis was that insurance providers could save money by investing in crisis centers to provide follow-up calls as both a measure to prevent suicidal behavior as well as the subsequent need for additional inpatient or emergency department intervention (Richardson, Mark, McKeon, 2013).

THE USE OF CRISIS HOTLINES IN FOLLOW-UP

Crisis hotlines are uniquely positioned to provide follow up care – crisis hotlines:

- Provide free, 24-hour access to staff trained in suicide assessment and intervention
- Thoroughly assess for risk of suicide, provide support, offer referrals, develop a safety plan, and dispatch emergency intervention, if necessary
- Connect directly with local mobile crisis teams
- Avert unnecessary ED visits and better ensure needed ED visits
- Intervene when a caller is not willing or able to ensure his or her own safety

Crisis centers have been shown to reduce emotional distress and suicidal ideation in callers (Gould, Kalafat, Munfakh, & Kleinman, 2007; Kalafat, Gould, Munfakh, & Kleinman, 2007).

Preliminary findings from an evaluation of crisis center follow-up to suicidal callers found that 80% of participants perceived the follow-up calls as helping at least a little in stopping them from killing themselves, while more than half of interviewed callers said the follow-up intervention helped a lot in stopping them from killing themselves. Callers who received more follow-up calls perceived the follow-up intervention to be more effective. Callers also perceived the intervention as more effective when counselors engaged in the following activities: discussing social contacts/settings as distractors; discussing social contacts to call when needing help; discussing warning signs; and exploring reasons for dying (Gould et al., unpublished).

Many Lifeline centers have a relationship with an ED and/or inpatient facility in their area.

- In Sacramento, California, WellSpace Health partners with Sutter Medical Center to provide follow-up for discharged patients. From 2010-2011, 98 patients consented to follow-up services and 93 were contacted by WellSpace Health. Of these, 90% were connected to referrals, 100% reported reduced distress, and 100% followed a safety plan. Only 2 re-attempts and readmissions were reported.
- In Cleveland, Ohio, FrontLine Service partners with 3 local hospitals. From 2010-2011, 49 patients consented to follow-up services and 46 were contacted by FrontLine Service. Of these, 100% reported lower risk for suicide, 72% followed a safety plan, and 50% linked to services referred by the crisis center. There were no reported attempts or readmissions.
- Thirteen Lifeline centers received SAMHSA funding in 2013 to undertake follow-up those discharged from local EDs and/or inpatient settings.

References:

For full list of references, please contact Naomi Carey at ncarey@mhaofnyc.org